

Supplementary Material for “Estimating the health benefit of
reducing indoor air pollution in a randomized environmental
intervention”

Roger D. Peng¹ Arlene M. Butz² Amber J. Hackstadt¹ D’Ann L. Williams³
Gregory B. Diette⁴ Patrick N. Breyse³ Elizabeth C. Matsui⁵

¹Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health

²Division of General Pediatrics, Johns Hopkins School of Medicine

³Department of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health

⁴Division of Pulmonary and Critical Care Medicine, Johns Hopkins School of Medicine

⁵Division of Pediatric Allergy and Immunology, Johns Hopkins School of Medicine

Corresponding Author: Roger D. Peng (rpeng@jhsph.edu)

Department of Biostatistics

Johns Hopkins Bloomberg School of Public Health

615 North Wolfe Street E3527

Baltimore MD 21205

USA

Phone: (410) 955-2468

Fax: (410) 955-0958




Figure S 1: Principal stratum treatment effects (posterior mean and 95% posterior intervals) in the original analysis (far right of each panel) and using a t -distribution with 2.5, 3, and 8 degrees of freedom to model in the (C) cleaner-under-treatment, (A) always-reduced, and (N) never-reduced groups.




Figure S 2: Principal stratum treatment effects (posterior mean and 95% posterior intervals) in the original analysis and with multiple imputation for the (C) cleaner-under-treatment, (A) always-reduced, and (N) never-reduced groups.